

To help lower blood sugar in adults with **type 2 diabetes** along with diet and exercise who are not at goal with oral diabetes medications

TALKING TO YOUR DOCTOR ABOUT YOUR A1C

You've been taking BYDUREON® BCise™ (exenatide extended-release) for injectable suspension for a few weeks now and soon you may be getting lab work done and discussing the results with your doctor. As he or she may have told you, your A1C test result reflects your average blood sugar level over the last 2 to 3 months, and it serves as a measure of how well you're managing your blood sugar.

The questions in this guide can help you have a productive conversation with your doctor about your A1C and how your treatment plan is progressing.

Questions You May Want to Ask Your Doctor:

- Can you help me understand what A1C is?
- What is my current A1C?
- What was my A1C from the last test?
- How has my A1C improved since my last visit?
- Is my A1C where it should be (at goal)? If not, what should my goal be?
- Are there any lifestyle changes I should consider that may help me get to goal?

IMPORTANT SAFETY INFORMATION

BYDUREON BCise may cause serious side effects, including:

- **POSSIBLE THYROID TUMORS, INCLUDING CANCER** Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In animal studies, BYDUREON and medicines that work like it caused thyroid tumors, including thyroid cancer. It is not known if BYDUREON BCise will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people
- **Do not use BYDUREON BCise if you or any of your family members have ever had MTC or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2)**
- **Do not use BYDUREON BCise if you have had an allergic reaction to exenatide** or any of the other ingredients in BYDUREON BCise

Please see additional Important Safety Information on the next page.

IMPORTANT SAFETY INFORMATION (cont'd)

- **Inflammation of the pancreas (pancreatitis)** Stop using BYDUREON BCise and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back
- **Low blood sugar (hypoglycemia)** Your risk for getting low blood sugar may be higher if you use BYDUREON BCise with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin. Tell your healthcare provider if you take other diabetes medicines. Talk to your healthcare provider about managing low blood sugar. **Signs and symptoms of low blood sugar include** dizziness or lightheadedness, sweating, confusion or drowsiness, headache, blurred vision, slurred speech, shakiness, fast heartbeat, anxiety, irritability, mood changes, hunger, weakness, or feeling jittery
- **Kidney problems (kidney failure)** Tell your healthcare provider if you have or had kidney problems. In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration) which may cause kidney problems to get worse
- **Stomach problems** Tell your healthcare provider if you have severe stomach problems, such as slowed emptying of your stomach (gastroparesis) or problems digesting food. Other medicines like BYDUREON BCise may cause severe stomach problems. It is not known if BYDUREON BCise causes or worsens stomach problems
- **Serious allergic reactions** Stop using BYDUREON BCise and get medical help right away if you have any symptoms of a serious allergic reaction, including itching, rash, or difficulty breathing
- **Injection-site reactions** Serious injection-site reactions, with or without bumps (nodules), have happened in some people who use BYDUREON. Some of these injection-site reactions have required surgery. Call your healthcare provider if you have any symptoms of injection-site reactions, including severe pain, swelling, blisters, an open wound, or a dark scab

The most common side effects with BYDUREON BCise may include a bump at the injection site and/or nausea. Nausea is most common when you first start using BYDUREON BCise, but decreases over time in most people as their body gets used to the medicine.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements, as taking them with BYDUREON may affect how each medicine works.

Before using BYDUREON BCise, tell your healthcare provider if you are pregnant, nursing, or plan to become pregnant or nurse. BYDUREON BCise may harm your unborn baby.

APPROVED USES

- BYDUREON BCise is an injectable prescription medicine that may improve blood sugar (glucose) in adults with type 2 diabetes mellitus, and should be used along with diet and exercise
- BYDUREON BCise is not recommended as the first choice of medicine for treating diabetes
- BYDUREON BCise is not a substitute for insulin and is not for people with type 1 diabetes or people with diabetic ketoacidosis
- Do not use BYDUREON, BYDUREON BCise, BYETTA or other medicines containing exenatide together
- It is not known if BYDUREON BCise can be used in people with a history of pancreatitis

Visit www.BYDUREONBCiseMedGuide.com for Medication Guide, www.BYDUREONBCise.com for Instructions for Use, and www.BYDUREONBCisePI.com for full Prescribing Information for BYDUREON BCise, including Boxed WARNING.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Once-weekly 
BYDUREON[®] BCise[™]
exenatide extended-release
injectable suspension 2 mg