

To help lower blood sugar in adults with **type 2 diabetes** along with diet and exercise who are not at goal with oral diabetes medications

READY TO TALK TO YOUR DOCTOR ABOUT PRESCRIPTION BYDUREON BCise?

Use this helpful guide

BYDUREON® BCise™ (exenatide extended-release) for injectable suspension is a once-weekly, non-insulin injection used along with diet and exercise that may help improve blood sugar (glucose) in adults with type 2 diabetes. BYDUREON BCise is not recommended as the first choice of medicine for treating type 2 diabetes.

This guide can help you have a productive conversation with your doctor about your type 2 diabetes and BYDUREON BCise. Please print and bring these questions with you to your next doctor appointment.

FIRST: Where am I now?

1. What is my current blood sugar (A1C) level?
2. Is my blood sugar (A1C) level where it should be?
3. If not, what should be my blood sugar goal?
4. In addition to my blood sugar level, are my current diet and exercise habits adequate?

NEXT: What next steps can help me reach my goals?

1. Is it time to change my treatment plan?
2. What lifestyle changes can I make that might help me get to my goal?
3. What additional treatment options are available for me?

FINALLY: Is once-weekly BYDUREON BCise a good option for me?

1. Could BYDUREON BCise, when used in addition to diet and exercise, help me lower my blood sugar?
2. Does BYDUREON BCise contain insulin?
3. How does one dose of BYDUREON BCise last for a whole week?
4. How does the unique delivery system in BYDUREON BCise make it different from other medicines?
5. How long does it take for BYDUREON BCise to work?
6. What should I know before taking BYDUREON BCise?
7. What do I need to know about injecting myself?

IMPORTANT SAFETY INFORMATION

BYDUREON BCise may cause serious side effects, including:

- **POSSIBLE THYROID TUMORS, INCLUDING CANCER** Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In animal studies, BYDUREON and medicines that work like it caused thyroid tumors, including thyroid cancer. It is not known if BYDUREON BCise will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people
- **Do not use BYDUREON BCise if you or any of your family members have ever had MTC or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2)**
- **Do not use BYDUREON BCise if you have had an allergic reaction to exenatide** or any of the other ingredients in BYDUREON BCise

Please see additional Important Safety Information on the next page.

Once-weekly 
BYDUREON® BCise™
exenatide extended-release
injectable suspension 2 mg

IMPORTANT SAFETY INFORMATION (cont'd)

- **Inflammation of the pancreas (pancreatitis)** Stop using BYDUREON BCise and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back
- **Low blood sugar (hypoglycemia)** Your risk for getting low blood sugar may be higher if you use BYDUREON BCise with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin. Tell your healthcare provider if you take other diabetes medicines. Talk to your healthcare provider about managing low blood sugar. **Signs and symptoms of low blood sugar include** dizziness or lightheadedness, sweating, confusion or drowsiness, headache, blurred vision, slurred speech, shakiness, fast heartbeat, anxiety, irritability, mood changes, hunger, weakness, or feeling jittery
- **Kidney problems (kidney failure)** Tell your healthcare provider if you have or had kidney problems. In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration) which may cause kidney problems to get worse
- **Stomach problems** Tell your healthcare provider if you have severe stomach problems, such as slowed emptying of your stomach (gastroparesis) or problems digesting food. Other medicines like BYDUREON BCise may cause severe stomach problems. It is not known if BYDUREON BCise causes or worsens stomach problems
- **Serious allergic reactions** Stop using BYDUREON BCise and get medical help right away if you have any symptoms of a serious allergic reaction, including itching, rash, or difficulty breathing
- **Injection-site reactions** Serious injection-site reactions, with or without bumps (nodules), have happened in some people who use BYDUREON. Some of these injection-site reactions have required surgery. Call your healthcare provider if you have any symptoms of injection-site reactions, including severe pain, swelling, blisters, an open wound, or a dark scab

The most common side effects with BYDUREON BCise may include a bump at the injection site and/or nausea. Nausea is most common when you first start using BYDUREON BCise, but decreases over time in most people as their body gets used to the medicine.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements, as taking them with BYDUREON may affect how each medicine works.

Before using BYDUREON BCise, tell your healthcare provider if you are pregnant, nursing, or plan to become pregnant or nurse. BYDUREON BCise may harm your unborn baby.

APPROVED USES

- BYDUREON BCise is an injectable prescription medicine that may improve blood sugar (glucose) in adults with type 2 diabetes mellitus, and should be used along with diet and exercise
- BYDUREON BCise is not recommended as the first choice of medicine for treating diabetes
- BYDUREON BCise is not a substitute for insulin and is not for people with type 1 diabetes or people with diabetic ketoacidosis
- Do not use BYDUREON, BYDUREON BCise, BYETTA or other medicines containing exenatide together
- It is not known if BYDUREON BCise can be used in people with a history of pancreatitis

Visit www.BYDUREONBCiseMedGuide.com for Medication Guide, www.BYDUREONBCise.com for Instructions for Use, and www.BYDUREONBCisePI.com for full Prescribing Information for BYDUREON BCise, including Boxed WARNING.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Once-weekly 
BYDUREON[®] BCise[™]
exenatide extended-release
injectable suspension 2 mg